Application of the Leray-Schauder Nonlinear Alternative

Břetislav Fajmon, Zdeněk Šmarda

Faculty of Electrical Engineering and Communication
VUT Brno, Technická 8, Czech Republic
e-mail: smarda@feec.vutbr.cz, fajmon@feec.vutbr.cz

Consider the following initial value problem

$$y'(t) = f(t, y(t), Ky(t)), \quad y(0) = y_0 \quad a.e. \quad t \in I, \tag{1}$$

where $I = [0, T]$, $f : I \times R^n \times R^n \to R^n$, and $Ky(t) = \int_0^T \kappa(t, s, y(s))ds$ is a Fredholm operator with $\kappa : I \times I \times R^n \to R^n$.

Similarly problems have been investigated by means of Schauder fixed point theorem or Schauder-Tychonoff fixed point theorem (see [1,2,3]).
Lerray-Schauder nonlinear alternative is an another approach to investigations of these ones.

Throughout this contribution the space of continuous functions on the interval I is denoted by C(I) with the norm $|.|_0$ given by

$$|y|_0 = \sup_{t \in [0,T]} |y(t)|.$$

$L^p(I)$, $1 \leq p \leq \infty$, is the space of measurable functions. For $y \in L^p(I)$the norm is given by

$$||y||_p = \left(\int_I |y(t)|^p dt \right)^{\frac{1}{p}} \ for \ 1 \leq p < \infty,$$

$$||y||_p = ess \ \sup_{t \in I} |y(t)| \ for \ p = \infty.$$

Finally, AC(I) is the space of absolutely continuous functions on I.
To obtain an existence theorem for (1) there is used Leray-Schauder Nonlinear Alternative type:

Le C be a convex subset of a normed linear space E, and let U be an open subset of C with $p^* \in U$. Then every compact, continuous map $\tilde{N} : \overline{U} \to C$ has at least one of the following two properties:

(I) $\tilde{N}$ has a fixed point.

(II) there is a $x \in \partial U$, with $x = (1 - \lambda)p^* + \lambda \tilde{N} x$ for some $0 < \lambda < 1$.

Definition 1. *A function $f : I \times R^n \times R^n \to R^n$, is L^1 -Carathéodory, if the following conditions hold:*

(i) the map $t \mapsto f(t,x,y)$ is measurable for all $(x,y) \in R^{2n}$,

(ii) the map $(x,y) \mapsto f(t,x,y)$ is continuous for a.e. $t \in I$,

(iii) for any $r > 0$, there exists $h_r \in L^1(I)$, such that, $|f(t,x,y)| \leq h_r(t)$, for a.e. $t \in I$, and for all $|x| \leq r$, $|y| \leq r$.

Definition 2: *A function $\kappa : I \times I \times R^n \mapsto R^n$ is integrable bounded L^1 -Carathéodory in t , if the following conditions hold:*

(i) the map $s \mapsto \kappa_t(s,y) = \kappa(t,s,y)$ is measurable for all $y \in R^n$,

(ii) the map $y \mapsto \kappa_t(s,y)$ is continuous for a.e. $s \in I$,

(iii) for any $r > 0$, there exists $\mu_{r,t} \in L^1(I)$, such that , $|y| \leq r$ implies $|\kappa_t(s,y)| \leq \mu_{r,t}(s)$ for a.e. $s \in I$,

(iv) $\sup_{t \in I} \int_I \mu_{r,t}(s)ds < \infty$.

Theorem. *Suppose the following conditions are satisfied :*

(i) the map $(t,s) \mapsto \kappa(t,s,x)$ is measurable for all $x \in R^n$, and the map $x \mapsto \kappa(t,s,y)$ is continuous for a.e. $(t,s) \in I \times I$,

(ii) κ is integrably bounded L^1 -Carathéodory in t,

(iii) f is L^1-Carathéodory .

In addition, suppose there exists a constant $M > 0$, independent of λ , with $|y_0| < M$, $|y|_0 \neq M$ for any solution $y \in AC(I)$ of

$$y'(t) = \lambda f(t, y(t), Ky(t)), \quad y(0) = y_0 \quad a.e.\ t \in [0,T],$$

for each $\lambda \in (0,1)$. Then (1) has at least one solution $y \in AC(I)$.

Acknowledgement

This research has been supported by the Czech Ministry of Education in the frame of MSM002160503 Research Intention MIKROSYN New Trends in Microelectronic Systems and Nanotechnologies.

References

[1] Agarwal,R.P., O'Regan,D.: *Singular Differential and Integral Equations with Applications*, Kluwer Academic Publishers, 2003, ISBN1-4020-1457-0.

[2] Meehan M., O'Regan,D.: *Existence theory for nonlinear Volterra integrodifferential and integral equations*, Nonlinear Analysis , 31(1998)m 317-341.

[3] Pachpate,B.G.: *Applications of the Leray-Schauder alternative to some Volterra integral and integrodifferential equations,* Indian J. Pure Appl. Math., 26(1995), 1161-1168.